

**Pre-Prescription Visits:**

- Discuss PrEP use; Clarify misconceptions
- Perform following laboratory tests:
  - HIV Test
  - Metabolic panel
  - Urinalysis
  - Hepatitis A, B, and C serology
  - STI screening
  - Pregnancy test



After confirmation of negative HIV test:  
**Prescribe 30-day supply of PrEP**  
Follow up in 2 weeks to assess side effects



**Adherence** and commitment should be assessed at **each visit**. Schedule visits every 30 days for patients who report poor adherence or intermittent use.



**30-day visit:**

Assess:

- Side effects
- Serum creatinine and calculated creatinine clearance for patients with borderline renal function or at increased risk for kidney disease (>65 years of age, black race, hypertension, or diabetes)
- Discuss risk reduction and provide condoms

**Prescribe 60-day refill; patient must come in for 3-month visit for HIV test and follow-up assessments, then 90-day schedule can begin.**

**3-month visit**

- HIV
- Ask about STI symptoms
- Discuss risk reduction and provide condom
- Serum creatinine and calculated creatinine clearance
- Pregnancy test

**6-month visit**

- HIV test
- Obtain STI screening tests
- Pregnancy test
- Discuss risk reduction and provide condoms

**9-month visit**

- HIV test
- Ask about STI symptoms
- Discuss risk reduction and provide condoms
- Serum creatinine and calculated creatinine clearance
- Pregnancy test

**12-month visit**

- HIV test
- Obtain STI screening tests
- HCV serology for MSM, IDUs and those with multiple sexual partners
- Discuss risk reduction and provide condoms
- Pregnancy test
- Urinalysis

